

# Concession Stand Tips

## SAFETY FIRST

### Requirement 9

#### 12 Steps to Safe and Sanitary

#### Food Service Events: The

#### following information is

#### intended to help you run a

#### healthful concession stand.

#### Following these simple

#### guidelines will help minimize

#### the risk of foodborne illness.

#### This information was provided

#### by District Administrator

#### George Glick, and is excerpted

#### from "Food Safety Hints" by

#### the Fort Wayne-Allen County,

#### Ind., Department of Health.

##### 1. Menu.

Keep your menu simple, and keep potentially hazardous foods (meats, eggs, dairy products, protein salads, cut fruits and vegetables, etc.) to a minimum. Avoid using precooked foods or leftovers. Use only foods from approved sources, avoiding foods that have been prepared at home. Complete control over your food, from source to service, is the key to safe, sanitary food service.

##### 2. Cooking.

Use a food thermometer to check on cooking and holding temperatures of potentially hazardous foods. All potentially hazardous foods should be kept at 41° F or below (if cold) or 140° F or above (if hot). Ground beef and ground pork products should be cooked to an internal temperature of 155° F, poultry parts should be cooked to 165° F. Most foodborne illnesses from temporary events can be traced back to lapses in temperature control.

##### 3. Reheating.

Rapidly reheat potentially hazardous foods to 165° F. Do not attempt to heat foods in crock pots, steam tables, over sterno units or other holding devices.

Slow-cooking mechanisms may activate bacteria and never reach killing temperatures.

##### 4. Cooling and Cold Storage.

Foods that require refrigeration must be cooled to 41° F as quickly as possible and held at that temperature until ready to serve. To cool foods down quickly, use an ice water bath (60% ice to 40% water), stirring the product frequently, or place the food in shallow pans no more than 4 inches in depth and refrigerate. Pans should not be stored one atop the other and lids should be off or ajar until the food is completely cooled. Check temperature periodically to see if the food is cooling properly. Allowing hazardous foods to remain unrefrigerated for too long has been the number ONE cause of foodborne illness.

##### 5. Hand Washing.

Frequent and thorough hand washing remains the first line of defense in preventing foodborne disease. The use of disposable gloves can provide an additional barrier to contamination, but they are no substitute for hand washing!

##### 6. Health and Hygiene.

Only healthy workers should prepare and serve food. Anyone who shows symptoms of disease (cramps, nausea, fever, vomiting, diarrhea, jaundice, etc.) or who has open sores or infected cuts on the hands should not be allowed in the food concession area. Workers should wear clean outer garments and should not smoke in the concession area. The use of hair restraints is recommended to prevent hair ending up in food products.

##### 7. Food Handling.

Avoid hand contact with raw, ready-to-eat foods and food contact surfaces. Use an acceptable dispensing utensil

to serve food. Touching food with bare hands can transfer germs to food.

##### 8. Dishwashing.

Use disposable utensils for food service. Keep your hands away from food contact surfaces, and never reuse disposable dishware. Wash in a four-step process:

1. Washing in hot soapy water;
2. Rinsing in clean water;
3. Chemical or heat sanitizing; and
4. Air drying.

##### 9. Ice.

Ice used to cool cans/bottles should not be used in cup beverages and should be stored separately. Use a scoop to dispense ice; never use the hands. Ice can become contaminated with bacteria and viruses and cause foodborne illness.

##### 10. Wiping Cloths.

Rinse and store your wiping cloths in a bucket of sanitizer (example: 1 gallon of water and 1/2 teaspoon of chlorine bleach). Change the solution every two hours. Well sanitized work surfaces prevent cross-contamination and discourage flies.

##### 11. Insect Control and Waste.

Keep foods covered to protect them from insects. Store pesticides away from foods. Place garbage and paper wastes in a refuse container with a tight-fitting lid. Dispose of wastewater in an approved method (do not dump it outside). All water used should be potable water from an approved source.

##### 12. Food Storage and Cleanliness.

Keep foods stored off the floor at least six inches. After your event is finished, clean the concession area and discard unusable food.

##### 13. Set a Minimum Worker Age.

Leagues should set a minimum age for workers or to be in the stand; in many states this is 16 or 18, due to potential hazards with various equipment.

*Safety plans must be postmarked no later than May 1st.*

# Volunteers Must Wash Hands

## HOW



## WHEN

**Wash your hands before you prepare food or as often as needed.**

### Wash after you:

- ▶ use the toilet
- ▶ touch uncooked meat, poultry, fish or eggs or other potentially hazardous foods
- ▶ interrupt working with food (such as answering the phone, opening a door or drawer)
- ▶ eat, smoke or chew gum
- ▶ touch soiled plates, utensils or equipment
- ▶ take out trash
- ▶ touch your nose, mouth, or any part of your body
- ▶ sneeze or cough

**Do not touch ready-to-eat foods with your bare hands.**

Use gloves, tongs, deli tissue or other serving utensils.  
Remove all jewelry, nail polish or false nails unless you wear gloves.

### Wear gloves.

when you have a cut or sore on your hand  
when you can't remove your jewelry

### If you wear gloves:

- ▶ wash your hands before you put on new gloves

### Change them:

- ▶ as often as you wash your hands
- ▶ when they are torn or soiled

Developed by UMass Extension Nutrition Education Program with support from U.S. Food & Drug Administration in cooperation with the MA Partnership for Food Safety Education, United States Department of Agriculture Cooperating. UMass Extension provides equal opportunity in programs and employment.

